

Bee Diseases

Braula

(This is a currently non - notifiable disease)

Braula (Braula coeca), also known as the beelouse, is found in colonies of the honey bee (*Apis mellifera*), where it lives on the bodies of the bees and literally steals its food out of the mouth of its host. Having said that it is neither a mite, a parasite, or true louse, but a tiny wingless fly. The beelouse is found on worker, drone, and queen bees, and as many as 187 have been recorded on a single queen bee!

It is generally believed that the beelouse causes little or no harm to bee colonies, but in beekeepers' apiaries they should be considered a pest as the larvae cause damage to the appearance of comb honey by burrowing under the cappings. Most beekeepers practice mechanical control of *Braula* unknowingly by extracting honey because the *Braula* larvae are eliminated while removing cappings before extraction. *Braula* also over-winter with adult the bees

Note: Because *Braula* larvae burrow under cappings it is advisable to freeze cut comb for a couple of days at -15C to ensure larvae and eggs are killed off.

Recognition:



Braula is a six legged insect or more precisely a wingless fly . It is remarkably similar to the Varroa mite (an arachnid with eight legs) at first glance due to its colouration, size and form, but on closer inspection the similarities disappear. Compare the two images on the left. The *Braula coeca* in the left image looks more spider-like than the flatter crab-like form of the Varroa mite on the right.

Life Cycle

Eggs: Eggs are white, oval-shaped, with two lateral flanges which are flat and extend parallel to each other and to the long axis of the egg. Including the flanges, a typical egg measured 0.84 mm by 0.42 mm. Eggs may be deposited in various places in a hive--in empty cells, on brood cell cappings, or on wax dirt on the floor of a colony, but only eggs that are laid on honey cappings hatch. Incubation period for eggs varies from two days during summer to 7.4 days during winter.

Larvae: Larvae emerge from the attached end of their egg where they begin constructing a tunnel under cappings and sometimes on the walls and bottoms of cells. These tunnels give cappings of infested comb the appearance of being intersected with fine fractures, similar to the mines of a leaf miner. Larvae feed upon honey and pollen grains within the wax of their tunnels. Bee louse larvae pass through three instars, requiring from 7.1 to 10.8 days to complete larval development, depending upon the season of the year.

Pupa: The prepupa ranges from 1 to 2.7 days and appears cream-white when viewed through the transparent larval skin. The pupa is enclosed within the unmodified cuticle of the last larval instar and is white or yellowish in color, 1.4 to 1.7 mm long by 0.5 to 0.75 mm wide.

Adult: The beelouse has eye rudiments just above the antennae seen as pale spots on the cuticle surface surrounded by more darkly pigmented chitinous rings. There is no trace of

wings or halteres. The tarsi are 5-segmented; each terminal joint contains a comb-like structure, divided in the middle, with a variable number of teeth. The combs allow Braula to cling tightly to the host.

Vectors

Braula can exist in beekeeping equipment, honey, wax, floor debris etc, and can spread quickly during normal hive / colony manipulations.

- Beekeepers - Transferring contaminated equipment / material between hives, colonies and apiary sites.
- Robbing - Weakened colonies will fall prey to robbing, transferring disease to other colonies and apiaries.
- Drifting - As with Robbing will transfer spores to other colonies.
- Swarming - Swarms can carry the lice with them to new sites where the disease will spread.

Note: beekeepers are the principal and most rapid means of spreading bee diseases.

Effects on Apis mellifera colonies

Not recognised a danger to honeybee colonies however it does cause damage to honey comb cell cappings.

How to Manage Braula

Braula (*Braula coeca*), also known as the bee louse, is found in colonies of the honey bee (*Apis mellifera*), where it lives on the bodies of the bees and literally steals its food out of the mouth of its host. Not generally believed to cause harm to bee colonies, but in beekeepers' apiaries they may be considered a pest as the larvae cause damage to the appearance of comb honey by burrowing under the cappings.

Detection

Hive Examination - Close examination of bees in the hive using a magnifying glass is the only way of identifying Braula presence on their hosts.

Monitoring

Vigilance is important with all honeybee diseases. Check all apiaries and colonies regularly for health and suspect any colonies that are not thriving where there is no already known reason. Colonies that die out should be examined thoroughly and sealed to prevent robbing and spread of any disease present.

Controls

There is no specific treatment recommended for Braula although it is suspected that Varroa treatments have the same lethal effect on Braula as on Varroa mites. The disease has negligible effect on healthy colonies. Prevention is the best method of controlling this disease by maintaining healthy, strong and vigorous colonies that display good hygienic traits. Good husbandry contributes greatly to overall colony behaviour and health, thereby avoiding the conditions in which disease can flourish.

Note: Its worthy of note that the bee louse lives naturally with honeybees as do other lice with their hosts. It is entirely possible that through other disease treatments the bee louse could become an 'endangered species' itself.

Apiary Housekeeping:

- Always maintain a high level of hygiene in all your beekeeping practices
- Carry out methodical health inspections on a regular basis, checking for brood disease particularly in spring and autumn.
- Never transfer combs between colonies without checking for brood diseases
- Systematically replace old brood combs in your hives melting down the old comb to maintain clean and healthy brood.
- Never bring colonies or equipment into your apiary without establishing their origin, condition, and disease status.
- Sterilise any secondhand equipment or hive components before introducing them into your apiary
- Discourage drifting and robbing in the apiary
- Suspect stray swarm health until you know otherwise
- Report any incidence of disease or suspicious conditions immediately to your local association